### 2026 | Volume Volume - 11 - Issue Issue - 2

#### In this issue

#### **Research Article**

Open Access Research Article PTZAID:ADA-11-202

# Anxiety and Depressive Symptoms in Patients with Hypothyroidism: A Cross-Sectional Study in Dakar, Senegal

Published On: November 05, 2025 | Pages: 025 - 031

Author(s): Momar Camara\*, Doriane Kaburundi, Sokhna Seck, Rokhaya Gueye, El Hadji Makhtar BA, Ngoné D Diack, Djiby Sow and Aïda Sylla

Background: Hypothyroidism is frequently associated with affective disturbances; however, the burden of anxiety and depressive symptoms among Senegalese patients remains poorly documented. Objective: This study aimed to determine the prevalence of anxiety and depressive symptoms and identify the associated clinical, biological, and sociodemographic factors among pati ...

Abstract View Full Article View DOI: 10.17352/2455-5460.000102

### **Review Article**

Open Access Review Article PTZAID:ADA-11-201

# Models of Depression for Preclinical Drug Discovery and Development: A **Transitional Perspective**

Published On: September 09, 2025 | Pages: 016 - 024

Author(s): Michel Bourin\*

Animal models of depression are most frequently encountered within the pharmaceutical industry to screen molecules and detect a putative antidepressant activity of a drug. The multiplicity of approaches and models makes comprehensive statements difficult, but animal models are necessary. These are experimental arrangements where a simple system is utilized to represen ...

Abstract View Full Article View DOI: 10.17352/2455-5460.000101

Open Access Review Article PTZAID:ADA-11-200

### **Evidence Psychotherapy Options for Borderline Personality Disorders**

Published On: August 19, 2025 | Pages: 008 - 015

Author(s): María-José Martín-Vázquez\*

Borderline Personality Disorder (BPD) is a prevalent and severe condition that affects emotional regulation, interpersonal relationships, and identity. BPD affects 1-2% of the population, mainly women, with high comorbidity, especially posttraumatic stress disorder, and a significant suicide rate. The diagnosis is based on ICD-11 and DSM-5 criteria with functional ev ...

Abstract View Full Article View DOI: 10.17352/2455-5460.000100